

DOI:

10.22301/IJHMCR.2528-3189.641

Article can be accessed online on:
<http://www.ijhmcr.com>

ORIGINAL ARTICLE

**INTERNATIONAL JOURNAL
OF HEALTH MEDICINE AND
CURRENT RESEARCH**

**TYPICAL CHARACTER IN THE SOUTH OF BANDA SEA
BASED ON THICKNESS AND VARIABILITY IN THE UPPER
LIMIT THERMOCLINE AREA AND ITS RELATIONSHIP
WITH SOUND VELOCITY**

**Hendry. I. Elim^{1,2,3,5,6,7,8}, Pieldrie Nanlohy^{1,2,3,8*}, Rayi Lalita^{1,2},
Nur Sahartira^{1,2}, Hanti Silawane^{1,2}, and Agus. S. Atmadipoera⁴**

¹ Physics Department, Faculty of Mathematics and Natural Sciences (FMIPA), Pattimura University (UNPATTI), Jl. Martinus Putuhena, Poka, Ambon, Indonesia 97233

² Computational Oceanography and Weather Laboratory (CO&W Lab.), Physics Department, Faculty of Mathematics and Natural Sciences (FMIPA), UNPATTI, Jl. Martinus Putuhena, Poka, Ambon, Indonesia 97233

³ Multidisciplinary Research Center of Excellence (MrCE), UNPATTI, Jl. Chr. Soplanit, RumahTiga, Ambon, Indonesia 97234

⁴ Department of Marine Science and Technology, FPIK, Institut Pertanian Bogor (IPB), Bogor, Jawa Barat, Indonesia

⁵ Nanomaterials for Photonics Nanotechnology Laboratory (Lab.N4PN),

⁶ Research Center of Nanotechnology and Innovative Creation (PPNRI-LEMLIT), UNPATTI, Jl. Chr. Soplanit, RumahTiga, Ambon, Indonesia 97234

⁷ Theoretical Physics Laboratory, Physics Department, FMIPA, UNPATTI Jl. Martinus Putuhena, Poka, Ambon, Indonesia 97233

⁸ Maritime and Marine Science Center of Excellence Jl. Dr. Leimena, Ambon, Indonesia 97234

ARTICLE INFO

Article History:

Received 07th November, 2017

Received in revised form

25th November, 2017

Accepted 01th December, 2017

Published online 23th December,
2017

Key words:

Thickness, Variability, Thermocline,
South of Banda sea, Sound velocity.

*Correspondence to Author:

Pieldrie Nanlohy

Physics Department, Faculty of
Mathematics and Natural Sciences
(FMIPA), Pattimura University

ABSTRACT

Typical character of south Banda Sea based on thickness and variability in the upper limit thermocline area has been identified by studying the whole area of Banda sea using the ~7 years INDESO data recorded from 3rd March 2007 to 18th March 2014. We found from such investigation of 4 types of seasons: (1) 2nd Middle Season (2ndMS), (2) West Season (WS), (3) 1st Middle Season (1stMS), and (4) East Season (ES) happened in the area that (1) 2ndMS, and (2) WS caused by North West Monsoon (NWM), and (3) 1stMS and (4) ES affected by South East Monsoon (SEM) contributed to the understanding of the upper layer of thermocline depth closely associated with the change of season. **The detail discussion of such finding especially related to sound velocity in each three different layers of mixed layer, thermocline, and below thermocline is provided in this present paper.**

Copyright © 2017, Pieldrie Nanlohy.. This is an open access article distributed under the creative commons attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Ocean is the most attractive field on earth due to its complex physical environmental system as well as its largest food contributions on earth which is about 70% in the total amount in comparison with that from the land area of the earth [1-4, 8-11]. Therefore, the environmental problem in the sea area can directly influence the life of human being on earth, of course with various complicated causes including its links with natural disasters [5-7, 12].

Banda sea is one of the earth deepest sea with its deep about 7000 m. Because of its deepest uniqueness, there was a lot of misteries beneath it particularly associated with various unique varieties of the ocean animals and plants living there. Banda area itself consisted of at least 11 islands. Based on the history of the islands, Maluku first official government was started there by Dutch due to spicies treasures in the islands. In order to understand the physical character in the Benda sea, there were many efforts conducted by a various multidisciplinary scientists such as biologists, geophysicists, marine scientists, and physicists. The contributions of such interdisciplinary scientists make the sea areas are interesting to be fully uncovered not only from foreigner scientists, but also from their tourists.

In this brief study, the authors present their research findings based on ~7 years data recorded from 2007 to 2014 of Infrastructure Development for Space Oceanography (INDESO) especially related to the typical character of south Banda Sea based on thickness and variability in the upper limit thermocline area as well as its connections with sound velocity in each three different layers of (1) Mixed layer (ML), (2) Thermocline layer (TL), and (3) Below Thermocline layer (BTL) as deep as ~318 m from the sea surface.

I. Computational Physics Method

The investigation method used in this current research was based on computational physics method particularly based on computational oceanography and weather in conjunction with computational and numerical algorithm analysis of a principal model in physical

system. The input data in this work was collected by INDESO in 7 years from 2007 to 2014. INDESO was a collaborative program between France and Indonesia. The row data had then been transformed into ordinary softwares such as Microsoft excel and origin to be analyzed and plotted. While by using Ocean Data View 4, the clear pictures of computational oceanography images can be highlighted, and monitored. In order to understand the scientific findings, ones needed to investigate the big pictures of Banda sea as well as all possibilities links with its thickness and variability influenced by the seasons with their typical weather as well as the sound velocity propagated in the 3 diferent upper layers of the sea such as (1) ML, (2) TL, and (3) BTL, respectively.

RESULTS AND DISCUSION

Figure 1 shows 4 types of seasons: (1) 2nd Middle Season (2ndMS), (2) West Season (WS), (3) 1st Middle Season (1stMS), and (4) East Season (ES) happended in the area that (1) 2ndMS, and (2) WS caused by North West Monsoon (NWM), and (3) 1stMS and (4) ES affected by South East Monsoon (SEM), respectively. **The significant change in Area 8 was due to WS caused by North West Monsoon (NWM).** While a little bit change of Area 8 affected by 1st MS was due to South East Monsoon (SEM).




Figure 1. 4 types of seasons: (1) 2nd Middle Season (2ndMS), (2) West Season (WS), (3) 1st Middle Season (1stMS), and (4) East Season (ES) happened in the area that (1) 2ndMS, and (2) WS caused by North West Monsoon (NWM), and (3) 1stMS and (4) ES affected by South East Monsoon (SEM), respectively.

Figure 2 depicts cyclonic Eddies effect of the improvement of the upper limit in the thermocline layer mainly happened because of WS happened in December to February located in the south part of Banda sea. The finding suggests that the upper limit of the thermocline layer in area (8) was the thinnest among other Banda sea areas.




Figure 2. Cyclonic Eddies effect of the improvement of the upper limit in the thermocline layer mainly happened because of West season (December to February) in the south part of Banda sea.

Figure 3 shows that November and January special behavior of sound velocity related to Monsoon and ENSO in Banda Sea have been identified using the data model from INDES0 recorded from July 2008 to May 2011, when ENSO phase was happened. From 9 areas of the whole Banda deep sea marked in the inset of Figure 1, we obtained such special physical phenomena in area no 3 called as North East of Banda area. Based on 3 types of sea vertical layers such as (1) ML, (2) TL, and (3) BTL on top of the deep Banda sea in such area, we observed that the ~4.54% top layers (≤ 318 m) of ~7000 m Banda deep sea shown a special character in November and January. It is interesting to note that only layers TL and BTL have a similar behavior. For example, in layer BTL, the El Nino phase changed the sound velocity to be weaker than the normal phase in July to May within 3 years from 2008 to 2011. On the other hand, the La Nina phase moved the sound velocity faster than the normal phase in July to April within the 3 observed years. Such phenomena was based on the observation data in this research. In our investigation significant finding as described well in Figure 3, the sound velocity is faster in the mixed layer than that in thermocline layer. While the below thermocline layer, we found that the sound velocity had the lowest speed.




Figure 3. Sound velocity propagation in 3 types of Banda sea layers such as mixed layers, thermocline layer, and below thermocline layer during 3 types of weather processes of Normal condition, El Nino and La Nina from July 2008 to May 2011, respectively.

Figure 4 describes that sound velocity was closely related with ASPL Nino 3.4. and wind velocity in which the speed of such sound decreases progressively from mixed layer to below thermocline layer. In the mixed layer of Banda sea, the sound

velocity has a significant fluctuation influenced by ASPL Nino 3.4. While in the thermocline layer of Banda sea, the wind velocity has a clear fluctuation due to a significant influence of such wind. As the speed of wind increases particularly from 6 to 8 m/s, the sound

velocity in the mixed layer slightly decreases from 1545 m/s to 1540 m/s.




Figure 4. (a) Sound velocity versus ASPL Nino 3.4, and (b) Sound velocity versus wind velocity in three layers (Mixed, Thermocline and Below Thermocline layers) of Banda Sea identified using the data model from Infrastructure Development for Space Oceanography (INDES0) recorded from July 2008 to May 2011, when ENSO phase was happened.

CONCLUSION

In conclusion, from 9 areas in the whole Banda deep sea, we obtained a unique physical phenomena in area no 3 called as North East of Banda area. Based on 3 types of sea vertical layers such as (1) ML, (2) TL, and (3) BTL on top of the deep Banda sea in such area, we observed that the **~4.54% top layers (≤ 318 m) of ~7000 m Banda deep sea shown a special character in November and January. This research suggests that such area no 3 of Banda sea may have a big source of many types of fish in the location.** In addition, the sound velocity in the mixed layer has an obvious fluctuation due to the speed of wind in the surface of the Banda sea. **While in the thermocline layer, the sound velocity was observed to have a great fluctuation in the low speed of wind of 1 to 5 m/s.**

ACKNOWLEDGEMENT

All the authors are grateful to thank the Department of Marine Science and Technology, **FPIK**, Institut Pertanian Bogor (**IPB**), Bogor, Jawa Barat, Indonesia for their financial supports in this research as well as Maritime and Marine Science Center of Excellence (**MMSCE**) for their research facilities.

REFERENCES

- [1]. Cole, J. E., 2000, Tropical Pacific Forcing of Decadal SST Variability in the Western Indian Ocean over the Past Two Centuries, *Science* Vol. 287(5453), pg. 617-619.
- [2]. Hakkinen, S., Rhines, P. B., and Worthen, D. L., 2011, Atmospheric Blocking and Atlantic Multi-decadal Ocean Variability, *Science* Vol. 334(6056), pg. 655-659.
- [3]. Deutsch, C., Brix, H., Ito, T., Frenzel, H., and Thompson, L.A., 2011, Climate-Forced Variability of Ocean Hypoxia, *Science* Vol. 333(6040) , pg. 336-339.
- [4]. Abram, N. J., Gagan, M. K., Cole, J. E., Hantoro, W. S., and Mudelsee, M, 2008, Recent intensification of tropical climate variability in the Indian Ocean, *Nature Geoscience* Vol. 1(12), pg. 849-853.
- [5]. Friedrich, T., Timmermann, A., Abe-Ouchi, A., Bates, N. R., Chikamoto, M. O., Church, M. J., Dore, J. E., Gledhill, D. K., González-Dávila, M., Heinemann, M., Ilyina, T., Jungclaus, J. H., McLeod, E., Mouchet, A., and Santana-Casiano, J. M., 2012, Detecting regional anthropogenic trends in ocean acidification against natural variability, *Nature Climate Change* Vol. 2(3), pg. 167-171.
- [6]. da Silva, A. E. and Véspoli de Carvalho, L. M., 2007, Large-scale index for South America Monsoon (LISAM), *Atmospheric Science Letters* Vol. 8(2), 51-57.
- [7]. Fleitmann, D., Burns, S. J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-Subbary, A.A., Buettner, A., Hippler, D., and Matter, A., 2007, Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra), *Quaternary Science Reviews* Vol. 26(1-2), pg. 170-188.
- [8]. Mizobata, K., Saitoh, S. I., Shiimoto, A., Miyamura, T., Shiga, N., Imai, K., Toratani, M., Kajiwara, Y., and Sasaoka, K., 2002 , Bering Sea cyclonic and

- anticyclonic eddies observed during summer 2000 and 2001, *Progress in Oceanography* Vol. 55 (1-2 SPEC ISS.), pg. 65-75.
- [9]. Gupta, A. K., Anderson, D. M., and Overpeck, J.T., 2003, Abrupt changes in the Asian southwest monsoon during the Holocene and their links to the North Atlantic Ocean, *Nature* Vol. 421 (6921n), pg. 354-357.
- [10]. Wang, B., and Ding, Q.H., 2006, Changes in global monsoon precipitation over the past 56 years, *Geophysical Research Letters* Vol. 33(6). DOI: 10.1029/2005GL025347.
- [11]. Shukla, J., 2007, Monsoon mysteries, *Science* Vol. 318, pg. 204-205.
- [12]. Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D'Arrigo, R. D., Jacoby, G. C., and Wright, W. E., 2010, Asian Monsoon Failure and Megadrought During the Last Millennium, *Science* Vol. 328(5977), pg. 486-489.
