

DOI:  
10.22301/IJHMCR.2528-3189.953

Article can be accessed online on:  
<http://www.ijhmcr.com>

-----  
ORIGINAL ARTICLE  
-----

INTERNATIONAL JOURNAL  
OF HEALTH MEDICINE AND  
CURRENT RESEARCH

## ANTIDIABETIC EFFECT OF INULIN FROM *Dioscorea esculenta* IN STREPTOZOTOCIN-INDUCED DIABETIC RATS

Anita Soraya Soetoko<sup>1</sup>, Azizah Hikma Safitri<sup>1</sup>, Israhanto Isradji<sup>1</sup>, Koko Agung Tri Wibowo<sup>1</sup>, Avicennia Nandira<sup>1</sup>, Dikta Zanwar Arifin Rafiq<sup>1</sup>, Salma Savita<sup>1</sup>, Ahmad N Al-Baari<sup>2</sup>, Yoyok Budi<sup>2</sup>, Dina Fatmawati<sup>1\*</sup>

<sup>1</sup> Faculty of Medicine, Universitas Islam Sultan Agung, Semarang, Indonesia.

<sup>2</sup> Faculty of Animal Husbandry and Agriculture, Universitas Diponegoro, Semarang, Indonesia.

### ARTICLE INFO

#### Article History:

Received 30th June, 2018

Received in revised form

25th July, 2018

Accepted 27th Agustus, 2018

Published online 30th September,  
2018

#### Key words:

Inulin, *Dioscorea  
esculenta*, antidiabetic effect.

#### \*Correspondence to Author:

**Dina Fatmawati**

Faculty of Medicine, Universitas  
Islam Sultan Agung, Semarang,  
Indonesia .

Jl. Raya Kaligawe KM.4, Semarang,  
Indonesia. Tel. +62246583584.

#### E-mail:

dienafatma@unissula.ac.id

### ABSTRACT

**Aims:** Antidiabetic effect of Inulin from *Dioscorea esculenta* tubers in streptozotocin-induced diabetic rats was investigated.

**Methods:** Antidiabetic effect was assessed based on HbA1C level, creatinine level, urea level, and microbiota composition in fecal. Level of HbA1C was determined using spectrophotometry. Level of creatinine and urea in blood serum was determined using enzymatic assay. The colony number of *Bifidobacterium sp* and *Lactobacillus sp* was evaluated using the total plate count method.

**Result:** After 14 days of treatment, there was a significant difference between control and inulin-treated groups in the level of HbA1C, creatinine, and urea. The colony numbers of *Bifidobacterium sp* and *Lactobacillus* were higher in inulin-treated groups than those of control group (p<0.05).

**Conclusion:** Inulin from *Dioscorea esculenta* tuber can be used as antidiabetic agent.

Copyright © 2018, Dina Fatmawati. This is an open access article distributed under the creative commons attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Citation:** Anita Soraya Soetoko<sup>1</sup>, Azizah Hikma Safitri<sup>1</sup>, Israhanto Isradji<sup>1</sup>, Koko Agung Tri Wibowo<sup>1</sup>, Avicennia Nandira<sup>1</sup>, Dikta Zanwar Arifin Rafiq<sup>1</sup>, Salma Savita<sup>1</sup>, Ahmad N Al-Baari<sup>2</sup>, Yoyok Budi<sup>2</sup>, Dina Fatmawati<sup>1\*</sup>, 2018 "Antidiabetic Effect Of Inulin From *Dioscorea esculenta* In Streptozotocin-Induced Diabetic Rats", *International Journal of Health Medicine and Current Research*, 3, (03), 953-957.

### INTRODUCTION

Diabetes mellitus is the most progressive metabolic disease associated with a constant increase in blood glucose level and impaired kidney function. Constant increased sugar levels may result in increased oxidative stress and production of glyco-oxidised products. Reactive oxidative species are

involved in pancreatic  $\beta$  cell dysfunction, apoptotic pathways, impaired insulin production, and insulin resistance<sup>1,2</sup>. Some studies suggest that type 1 diabetes mellitus is associated with changes in intestinal microbiota composition<sup>2</sup>. These changes occur along with the progression of diabetes leading to unfavorable intestinal microenvironment for certain types of bacteria<sup>3</sup>. Zhang *et al.*, (2013) showed that there was a decrease in the colony number of *Bifidobacterium sp* and *Lactobacillus sp* in patients with diabetes mellitus. The dysbiosis condition can be one of the cause in progression of diabetic nephropathy<sup>4</sup>.

Selective modulations of intestinal microbiota by prebiotics have been suggested to decrease inflammation and metabolic dysfunction in rats. Inulin is one of the prebiotics used to modulate intestinal microbiota<sup>5</sup>. Inulin can be isolated from tuber plants such as *Dioscorea esculenta*. *Dioscorea esculenta* is one type of tubers belonging to *Dioscoreaceae* family which is widely grown Indonesia. Previous studies showed that the *Dioscorea esculenta* tuber has anti-inflammatory, antioxidant and cytotoxicity effects on breast cancer cells<sup>6-8</sup>. Another study showed that *Dioscorea esculenta* has the highest inulin content (14.77% of dry weight) among the other *Dioscorea* family<sup>9</sup>.

Inulin fermentation has been shown to produce acetic acid of 137 $\pm$ 75 mmol, propionic acid of 11 $\pm$ 9 mmol and, butyric acid of 20 $\pm$ 17 mmol<sup>10</sup>. The products modulates levels of some intestinal hormones involved in glucose and energy homeostasis, including glucagon-like peptide (GLP) -1 and reduces blood glucose levels<sup>5</sup>. Inulin at a dose of 10 g/day has been preferred as a type of prebiotics due its tolerance and increase the number of *Bifidobacterium*<sup>11</sup>.

The purpose of this study was to investigate the effect of antidiabetic effect of inulin from *Dioscorea esculenta* based on HbA1C, creatinine and urea level as well as the colony number of *Bifidobacterium sp* and *Lactobacillus sp*.

## METHODS

All the research procedures have been approved by of Bioethical commission, Faculty of Medicine, Universitas Islam Sultan Agung (registration number: reg58/II/2018).

### Authentication of *Dioscorea esculenta*

*Dioscorea esculenta* whole plant was obtained from the area of Tugel Mountain, Purwokerto and was authenticated in the plant taxonomy laboratory, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang.

### Preparation of inulin powder from *Dioscorea esculenta*

*Dioscorea esculenta* powder was obtained by using cabinet drying methods. Fresh *Dioscorea*

*esculenta* was washed and sliced, then added with aquades with ratio of 1: 3, then blend evenly. The result was filtered using a filter cloth and stored in a refrigerator at temperature of -20°C for 24 hours. After 24 hours the frozen blend was thawed at room temperature and centrifuged at 4500 rpm for 10 minutes. The resulting precipitate was transferred into a petri dish and dried at 37°C for 24 hours. After drying, the precipitate was sieved into powder and then filtered.

### Concentration of Inulin Test

Concentration of inulin from *Dioscorea esculenta* was measured using carbazole cysteine method. 1 ml of filtrate of *Dioscorea esculenta* was added with 0.2 ml of cysteine 1.5% and 6 ml of 70% H<sub>2</sub>SO<sub>4</sub>. The mixture was then shaken, then added with 0.2 ml of 0.12% carbazole in an ethanol solution and heated at 60 ° C. for 10 minutes. After cold absorbance the sample was measured at a wavelength of 560 nm.

### Induction of diabetes in rats

The rats were fasted for 10 h before the intraperitoneal injection of streptozotocin (STZ) at the dose of 45mg/kg BW dissolved in 1 ml of 0.1 M-buffer citrate, pH 4.5. After 72 hours, blood glucose levels of the rats were evaluated using spectrophotometric method. The rats with fasting blood glucose range of above 135 mg/dl were considered as diabetic rats<sup>12</sup>.

### Experimental design

The treatment was conducted at Biology Laboratory, Faculty of Medicine, Universitas Islam Sultan Agung. A total of 10 male white Wistar rats were acclimatized for 7 days. During acclimatization, the rats were fed with a normal pellet diet and drinking water *ad libitum*. Then, 10 rats were divided into 2 as follows: P1: 5 STZ induced diabetic control fed with standard and aquades from day 5 to day 28, P2: 5 STZ diabetic rats administrated with inulin from from the *Dioscorea esculenta* tuber at the dose of 180 mg/rat/day from day 5 to day 28.

### Biochemical assays

HbA1C, Urea and creatine level were evaluated on day 29. Blood sample collected for estimation of HbA1C, creatinine, and urea was taken from the *retro-orbital plexus*, as much as 3 ml. Serum creatinine and urea serum levels were examined at Semarang Laboratory of Health Laboratory by enzymatic method was used to determine the serum creatinine and urea levels. Serum creatinine was measured using the Jaffe method without deproteinization, whereas serum urea levels were measured using the enzymatic Endpoint/Berthelot method. HbA1C levels was measured using the immunoassay and expressed in percentage.

### Bacterial species identification and quantitative colony count

Identification of *Bifidobacterium sp.* and *Lactobacillus sp.* was based on gram staining on culture of fecal bacteria grown on MRS medium, whereas the of bacterial colony counting was determined by total plate count method (TPC). The identification and quantity calculations were performed at Microbiology Laboratory, Faculty of Medicine, Universitas Islam Sultan Agung.

*Lactobacillus sp.* on MRS media was observed as flat edge (smooth), convex elevation, small colony shape, creamy color, and smooth shiny surface. Type *Bifidobacterium sp.* was identified by observe ng non-spore forming bacteria, positive gram characteristics, catalase negative, non-motile, pairs of V-shaped or Y-shaped pairs, in gram-positive staining. Colonies of *Bifidobacterium sp.* Was observed as a gray in color.

### Data analysis

Results were presented as mean (SD). Mean difference of HbA1C level of urea, creatinine, *Bifidobacterium sp* and *Lactobacillus sp* colony number between treatment group and control was analyzed using unpaired T test with 95% confidence interval.

### RESULTS

This study showed that inulin from *Dioscorea esculenta* lowered the level of of HbA1C, the creatinine, and urea in diabetic rats. Table 1 showed mean percentages of HbA1C, the creatinine, and urea level in treated group and diabetic control rats.


**Table 1.** Level of HbA1, creatinine and urea level in treated groups (mean SD).

Groups	HbA1C level in blood serum (%)	Creatinin level in blood serum (mg/dl)	Urea level in blood serum (mg/dl)
P1	7.62±0.50 <sup>a</sup>	0.81±0.06	83.17±14.8
P2	6.14±0.35 <sup>b</sup>	0.62±0.04	51.20±0.58

Note: the letters<sup>a, b</sup> which differ between treatment groups (P1 and P2) showed significant differences for each variable (p <0.05)

This study also showed that mean level of serum creatinine and urea in STZ-induced diabetic rats, in the experimental group was lower compared with that of the control group (p=0.001 vs p=0.006). The mechanism of decrease in creatinine

and urea levels after inulin administration was suspected to be associated with an increase in the number of *Bifidobacterium sp* and *Lactobacillus sp.* The colony number *Bifidobacterium sp* and *Lactobacillus sp* is presented in figure 1.



**Figure 2.** Number of *Bifidobacterium sp.* and *Lactobacillus sp.* colony (mean SD) (\*\*) Significant at p <0.05.

### DISCUSSION

There was a significant reduction in levels of HbA1C, creatinine and urea in inulin treated animals as compared with diabetic control. Heinemann and Freckmann (2015) have reported that the decrease in HbA1C levels is closely associated with the decrease in blood glucose levels, which can be estimated at 6.5%

(47.5 mmol/mol) HbA1C levels equivalent to blood glucose levels of 140 mg/dl, while HbA1C 8.6% (70.5 mmol/mol) is equivalent to a blood glucose level of 200 mg/dl<sup>13</sup>.

This present study showed that the inulin treatment significantly increase the number of *Bifidobacterium sp.* and *Lactobacillus sp* in the experimental group (p < 0.05) as compared to animal

control. This effect was likely due the capability of the inulin to stimulate the growth of bacteria *Bifidobacterium sp.* and *Lactobacillus sp.* as previously demonstrated by Winarti *et al.*, (2013) that the inulin from gembili (*Dioscorea esculenta*) can stimulate the growth of *Bifidobacterium breve* BRL-131, *Bifidobacterium bifidum* BRL-130, *Bifidobacterium longum* ATCC-15707 and *Lactobacillus acidophilus* in *invitro*<sup>14</sup>. This finding is also supported by the study of Pompei *et al.*, (2008) showing that inulin can increase the growth of *Bifidobacterium adolescentis*, *Bifidobacterium infantis*, *Bifidobacterium breve*, *Bifidobacterium longum*, *Lactobacillus plantarum*, *Lactobacillus rhamnosus*, *Lactobacillus reuteri*, *Lactobacillus delbruechi* and can inhibit the growth. *E. coli* and *Clostridium sp*<sup>15</sup>. The increase in the number of the two intestinal microbiota bacteria in our study might have been associated with the increased production of short chain fatty acids (SCFA) such as lactate, acetate, propionate and butyrate affecting glucose metabolism leading to increase production of inulin. This possible mechanism has been proposed by zhang *et al* (2013) in there study on the effect of the intestinal bacteria on the level of SCFA levels<sup>4</sup>. The increase of SCFA have been shown to increase insulin level via GLP-1<sup>16,5</sup>.

## CONCLUSION

Inulin from *Dioscorea esculenta* tuber has an antidiabetic effect in streptozotocin-induced rats by decreasing HbA1C, this may be due to a certain modulation effect on bacteria *Bifidobacterium sp.* and *Lactobacillus sp.* which is associated with decreased levels of creatinine and urea.

## ACKNOWLEDGMENTS

The Authors wish to thank Kemeristek DIKTI (contract number: 257/B.1/LPPM-SA/VI/2017), and Universitas Islam Sultan Agung for funding the research Drs. Eling Purwantoyo, M.Si. for his assistance in authentication of the *Dioscorea esculenta*.

## REFERENCES

1. Sharma P., Bhardhwaj P., Singh R.. (2016). Administration of Lactobacillus casei and Bifidobacterium bifidum Ameliorated Hyperglycemia, Dyslipidemia, and Oxidative Stress in Diabetic Rats. *Int J Prev Med.* 7:102.
2. Dominguez-Bello, M. G., Costello, E. K., Contreras, M., Magris, M., Hidalgo, G., Fierer, N. & Knight, R. (2010). Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci* 107, 11971–11975.
3. Patterson E, Marques TM, O’Sullivan O, Fitzgerald P, Fitzgerald GF, Cotter PD, Dinan TG, Cryan JF, Stanton C., Ross RP. Streptozotocin-induced type-1-diabetes disease onset in Sprague–Dawley rats is associated with an altered intestinal microbiota composition and decreased diversity. *Microbiology* (2015), 161, 182–193.
4. Zhang, X., Gregg, E. W., Williamson, D. F., Barker, L. E., Thomas, W., Bullard, K. M., ... & Albright, A. L. (2010). A1C level and future risk of diabetes: a systematic review. *Diabetes care*, 33(7), 1665-1673.
5. Cani, P. D., Geurts, L., Matamoros, S., Plovier, H., & Duparc, T. (2014). Glucose metabolism: focus on gut microbiota, the endocannabinoid system and beyond. *Diabetes & metabolism*, 40(4), 246-257.
6. Olayemi JO., Ajaiyeoba EO. (2007). Anti-inflammatory studies of yam (*Dioscorea esculenta*) extract on wistar rats. *African Journal of Biotechnology*. 6(16).
7. Murugan M., Mohan VR. (2012). In vitro antioxidant studies of *Dioscorea esculenta* (Lour). Burkill. *Asian Pacific Journal of Tropical Biomedicine* (S1620-S1624).
8. Fatmawati D., Mubarika S, Wahyuningsih, MSH. (2016). The effect of combined doxorubicin and *Dioscorea esculenta* (Lour.) Burk. on apoptosis induction in human breast cancer cell. *AIP Conference Proceedings* 1744 (1), 020011. <https://doi.org/10.1063/1.4953485>.
9. Winarti, S., Harmayani, E., & Nurismanto, R. (2011). Karakteristik dan profil inulin beberapa jenis uwi (*Dioscorea spp.*). *Agritech*, 31(4).
10. Boets E., Deroover L., Houben E., Vermeulen K., Gomand SV., Delcour JA., Verbeke K. 2015. Quantification of in Vivo Colonic Short Chain Fatty Acid Production from Inulin. *Nutrients*. vol 7(11) 8916-29. doi: 10.3390/nu7115440.
11. Mitchell CM, Davy BM, Halliday TM., Mathew W. Hulver MW., Andrew P. Neilson AP., Monica A. Ponder MA., Davya KP. (2015). The Effect of Prebiotic Supplementation with Inulin On Cardiometabolic Health: Rationale, Design, and Methods Of A Controlled Feeding Efficacy Trial in Adults at Risk of Type 2 Diabetes. *Contemp Clin Trials* 45(0 0): 328–337.
12. Puspitasari, A.P. and Syauqy, A., (2015). Pengaruh Pemberian Pisang Kepok (Musa Paradisiaca Forma Typical) Terhadap Kadar Malondialdehyde (Mda) Tikus Sprague Dawley Pra-Sindrom Metabolik. *Journal of Nutrition College*, 4(4), pp.314-322.
13. Heinemann L., Freckmann G. (2015). Quality of HbA1c Measurement in the Practice: The

- German perspective. *J Diabetes Sci Technol.* 9(3): 687–695.
14. Winarti, S; Harmayani, E; Marsono, Y; Pranoto, Y. (2013). Effect of inulin isolated from lesser yam (*Dioscorea esculenta*) on the growth of probiotics bacteria and SCFA formation during fermentation. *International Research Journal of Microbiology*, 4(2), 53–63.
15. Pompei A, Cordisco L, Raimondi S, Amaretti A, Pagnoni AM (2008). In vitro comparison of the prebiotic effect of two inulin-type fruktans. *Aerobe* 14: 280-286.
16. Byrne, C. S., Chambers, E. S., Morrison, D. J., & Frost, G. (2015). The role of short chain fatty acids in appetite regulation and energy homeostasis. *International journal of obesity*, 39(9), 1331.

\*\*\*\*\*