

**DOI:**

10.22301/IJHMCR.2528-3189.1703

Article can be accessed online on:

<http://www.ijhmcr.com>**ORIGINAL ARTICLE****INTERNATIONAL JOURNAL
OF HEALTH MEDICINE AND
CURRENT RESEARCH****COVID-19 PHYSIOLOGICAL AND MOLECULAR MECHANISM
RESULTING IN THREAT TO HUMAN SOCIETY-A REVIEW****K.R.Padma¹ K.R.Don² and P.Josthna³**^{1&3}Assistant Professor, Department of Biotechnology, Sri Padmavati Mahila VisvaVidyalayam (Women's) University, Tirupati, AP.²Reader, Department of Oral Pathology, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Velappanchavadi, Chennai, Tamil Nadu, India.**ARTICLE INFO****Article History:**Received 18th Jan, 2020,

Received in revised form

18th March, 2020Accepted 22th June, 2020Published online 30th June, 2020**Key words:**

SARS-CoV, COVID-19, Corona virus, Pandemic disease, Positive stranded RNA virus

Correspondence to Author:*Dr.K.R.Padma,**

Assistant Professor,

Department of Biotechnology,

Sri Padmavati Mahila Visva

Vidyalayam (Women's) University,

Tirupati, AP.

E-mail:

thulasipadi@gmail.com

ABSTRACT

Human Corona Viruses (HCoVs) are the displaying pandemic disease globally. It is commonly regarded as the pathogens of the respiratory system coupled in the midst of broad assortment of respiratory diseases. These HCoVs also revealed to possess an enveloped positive-sensed RNA viruses and multiplies mainly in the protoplasm of the host. It is been acknowledged that the virus common route of its entry is by contacting and later coalescing with the external surface of the host. Once entered the niceties of the corona virus need to be clearly understood. Although several advancements acknowledged about its structural properties as well as emphasized physicochemical and molecular mechanism of the coronavirus yet unable to distinctively comprehend on the unusual property of trimeric S protein. Characteristically the action of the coronavirus fusion peptide (FP) and induction of consensual alteration in the viral fusion proteins necessitate for comprehension. In our present review, we have imparted a concise introduction to corona viruses and discussed on their replication, physiology and molecular mechanism of pathogenicity, further portray different types of corona viruses. Hence this review noticeably highlights the outbursts of the mutated form of (SARS-CoV) commonly named as Severe Acute Respiratory Syndrome. Corona virus now basically regarded COVID-19 all over world.

Copyright © 2020, **Prof. Dr. Dipak Bhargava**. This is an open access article distributed under the creative commons attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: K.R.Padma¹ K.R.Don² and P.Josthna³, 2020 "Covid-19 Physiological And Molecular Mechanism Resulting In Threat To Human Society-A Review", International Journal Of Health Medicine And Current Research, 5, (02), 1703-1717.

1. Introduction

Approximately the last few months, globally there has been the subsistence rise of latest mutant forms of viruses that shroud insistent menace among worldwide well being. Lately, in December 2019 especially numerous patients from Wuhan, China were the first to exhibit conditions that were similar to flu as well as pneumonia and lead to outbursts throughout the world at a robust rate. Eventually, after understanding the severity of the cases the World Health Organization (WHO) acknowledged the growth of virus and arrived at a conclusion that this virus was quite different from SARS. Hence, renamed first as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) due to its similar symptoms [1]. Although, it belongs to Coronaviridae family but regarded as new virus as well as initially been acknowledged in the year 2019 so, regarded as novel coronavirus (2019-nCoV). Recently, the pathological diseases it aggravates lead to be recognized as COVID-19. Both the viruses COVID-19 as well as SARS-CoV are positive-sensed RNA virus originating from the family Coronaviridae [2-3].

In fact the strains belonging to coronavirus family reported in early 1930s [4]. Later the popularity of corona viruses acknowledged after the (SARS) sudden occurrence which juddered the humankind in 2002–2003. Repercussion of epidemic virus, interestingly hue again on this family of viruses and lead to the identification of COVID-19 belonging to these family members. At present the chaos it is causing and the enormity for society physical conditions in relation with diseases connected among humans and animals was raising interest. In fact this Corona viruses is not only persistent to humans but infects a vast variety of

animal creatures as well as avians, resulting in breathing problems as well as gastrocolic diseases and in few abnormal cases leads to liver cirrhosis as well as brain related diseases. Although the viral infection can result either in acute condition or continuous condition [5]. Based on the serological and genetic analysis, corona virus was recognized with four diverse subfamily such as α , β , γ , and δ -CoV [6].

All the four subfamily possess a protected spike protein which is anchored on the external surface as well as provides channel for the corona virus ingress into host cells [7]. Literally the protein spike known to be epitomized in different forms i.e basically the pre-fusion structure existing on developed virions along with post-fusion shape which is formed later blending along with the host outer membrane. Hence in our study we have focussed on the ingress strategies of corona viruses and unravel how these mechanisms are related to host tropism as well as pathogenicity. Since the virus is spreading globally, the WHO on March 11, 2020, officially described the COVID-19 outbreak as a pandemic in upcoming days and strategies needed to be undertaken to break the chain and manage the disease spreading.

2. Corona Virus Genome and domains of Spike Protein

The Corona viruses are pleiomorphic spherical, enveloped viruses, with varying dimensions from 80 to 120 nm. The 5' capped positive stranded RNA viral genome with was found to be approximately in the range of 27 and 34 kb, the largest in midst of all RNA viruses. The organization of the genome with approximately six to ten open reading frames (ORFs). Nevertheless the beginning ORF occupies major part of the genome and ascertain to posses the

replication proteins, while the final end possess the structural gene proteins arranged in a predetermined order: (HE)-S-E-M-N as shown in (Figure 1A). The genome is protected with envelope made of helical nucleocapsid surrounding the host-derived lipid bilayer. The three viral proteins were found on the outer virion envelope, one among them included the spike protein (S), the (M) protein referred as

Membrane protein and third enveloped protein (E) (Figure 1B). Additionally, few corona viruses encompassed hemagglutinin esterase (HE) along with membrane and enveloped proteins present on the envelope are essentially indulging in assembly of viruses, the S protein is the principal mediator for viral entry. [8-12].

Figure 1A




Figure-1B




Figure-1A & 1B: The genome organization of Corona virus and Structure of Corona encompassing diverse viral proteins

4. Inhibition of PERK pathway attenuates viral infection

Translation of proteins weakens the viral replication has been broadly monitored as a combat method for the host cells opposed to viral infection. Hence in order to reduce the transportation of viral proteins, replication of the virus would be hindered as well as proliferation of infection can also be restricted, thereby providing adequate time for the natural defenses to instigate effectual antiviral reactions. PERK pathway attenuates viral infections by eIF2 α kinases, thereby PKR results in Interferon-triggering reaction and precise identification of viral RNAs, plays a crucially significant task in triggering attenuation of translation in infected viral cells [32]. The novel viruses brilliantly have developed diverse methods to thwart PKR. Nevertheless, the non-structural 5A (NS5A) protein of Human corona virus unswervingly interacted with the active site of PKR, thus blocks PKR activation [33-37]. During the enhancement of virus infection, amass assembly of viral proteins can overwhelm blocks that have folding capacities for ER.

Earlier the S protein in SARS-CoV also has shown to activate ERK phosphorylation and revealed to augment IL8 release [38-45]. Furthermore, viral proteins of SARS-CoV have displayed in ERK triggering reaction. SARS-CoV Papain Like protease as well as SARS-CoV 3b expression proteins have shown significantly to enhance ERK1 ubiquitin-arbitrated disintegration to repress IFN-triggered reactions [46-49]. Although, SARS-CoV, MERS-CoV infection and now novel COVID-19 is also combined in triggering ERK phosphorylation setup. Nevertheless the ERK phosphorylation was not substantially influenced by HCoV-229E infection, since this mutated form is resistant to phosphorylated ERK but can be instigated by the use of Hydroxychloroquine a known antimalarial agent which was found to be effective against viruses [50]. Hence, profound comprehension on ERK pathway triggering method could reveal to be substantial competence in the period of HCoV infection [51].




Figure-3: Attenuation of Corona virus infection reduces translation of by PERK pathway

5. Emerging COVID-19 reports from Wuhan city in China including various countries

WHO reports of March 10, 2020, approximately 113,702 validated cases internationally, and 4012 mortalities have been recorded; approximately 75% of all authenticated cases (80,924) and 80% of overall demises are associated with COVID-19. Amongst,(3140) cases began from China and its territories. Regardless, the beginning of this virus cases came to notice from Wuhan city in China. Furthermore, WHO has formally announced China as a “Red alert zone” region for COVID-19 [\[52-54\]](#).

6. Chinese continent and surrounding areas

In China and its region of Hubei, involves Wuhan. Nevertheless, recently has been reported the largest authenticated COVID-19 positive cases and mortalities in areas of China (67,760 cases and 3,024 deaths). Amongst several regions in China has shown to cross the thousands mark especially in Guangdong, Henan, Zhejiang, and Hunan each have 1,353, 1,272, 1,215, and 1,018 cases, respectively during March beginning (Table -1). Later greater number of demises was registered in Guangdong with more than 22 cases [\[55-58\]](#).

Places	Confirmed cases	Deaths
Hubei	67,801	3160
Beijing	554	8
Liaoning	127	2
Anhui	990	6
Chongqing	578	6
Shaanxi	248	3
Tianjin	141	3
Fujian	318	1
Hunan	1018	4
Sichuan	545	3
Gansu	136	2
Heilongjiang	484	13
Jiangxi	936	1
Guangdong	1428	8
Shanghai	414	4
Yunnan	176	2
Zhejiang	1240	1

Table-1: Epidemiology of COVID-19 in Main land china (March

Table-2: COVID-19 reports of other countries

Country	Overall cases	Latest cases	Overall deaths	Latest deaths	Overall recovered	Positive cases	Danger state	Entire cases
Italy	63,927	-	6,077	-	7,432	50,418	3,204	1,057
USA	46,145	+2411	582	+29	295	45,268	1,040	139
Spain	35,136	-	2,311	-	3,355	29,470	2,355	751
Germany	29,056	-	123	-	453	28,480	23	347
Iran	23,049	-	1,812	-	8,376	12,861	-	274
France	19,856	-	860	-	2,200	16,796	2,082	304
South Korea	9,037	+76	120	+9	3,507	5,410	59	176
Switzerland	8,795	-	120	-	131	8,544	141	1,016
UK	6,650	-	335	-	135	6,180	20	98
Netherlands	4,749	-	213	-	2	4,534	435	277
Austria	4,477	+3	21	-	9	4,477	16	497
Belgium	3,743		88	-	401	3,254	322	323
Norway	2,625	-	10	-	6	2,609	41	484
Australia	2,136	+249	8	+1	118	2,010	11	84
Canada	2,091	-	24	-	320	1,747	1	55
Portugal	2,060	-	23	-	14	2,023	47	202
Sweden	2,046	-	27	-	16	2,003	104	203
Brazil	1,924	-	34	-	2	1,888	18	9

7. Emerging COVID-19 cases in INDIA

In India there are 30 states and Union territories which went under lock down as [HCoV](#) cases in India spiked to 471 on Monday 23/03/2020 as shown in (Table-3). Various places throughout India imposed

curfew in their own states. All domestic flights from midnight have been cancelled to refrain the virus spread, stepping up a nationwide lockdown. Even though the virus continues to spread worldwide, with the total number of infected cases reaching 378,741 and 16,502 death.

Table-3: COVID Data upto date

Name of State/UT	Confirmed cases (Indian)	Confirmed cases (Foreigners)	Deaths
Andhra Pradesh	7	0	0
Bihar	2	0	1
Chattisgarh	1	0	0
Delhi	30	1	1
Gujarat	29	0	1
Haryana	12	14	0
Himachal Pradesh	3	0	1
Karnataka	37	0	1
Kerala	87	8	0
Madhya Pradesh	7	0	0
Maharastra	84	3	2
Odisha	2	0	0
Puducherry	1	0	0
Punjab	21	0	1
Rajasthan	31	2	0
Tamil Nadu	10	2	0
Telangana	22	10	0
Union territory of Chandigarh	6	0	0
Union territory of J&K	4	0	0
Union territory of Ladakh	13	0	0
Uttar Pradesh	32	1	0
Uttarakhand	3	0	0
West Bengal	7	0	1
Total	451	41	9
Grand Total	492		




Figure-4 Cases of COVID-19 bar graph in India

8. Clinical signs and symptoms of COVID-19

The symptoms might be symptomatic or symptomless infections which have been described [59-63], but number of their frequency occurrence is unknown. As in cases with asymptomatic infection on chest computed tomography (CT), greater than 60 percent had displayed prototypical ground-glass opacities or speckled darkening imaging abnormalities [64]. In few cases arised low-levels of ailments, with or without other representative manifestations, are following few days on diagnosis.

9. Clinical manifestations

The preliminary clinical manifestation of the disease enhancement was pneumonia which happens to be the utmost regular staid materialization of infection, considered predominantly with ailment, tussis, hyperpnea and bilateral infiltrates on chest imaging [65-66]. At present stage of existence allow no particular clinical features that can comprehensibly and authentically differentiate COVID-19 from other viral respiratory diseases. At present scenario of Wuhan,

where outbreak of COVID-19 occurred described that 138 patients suffered from viral pneumonia in Wuhan, the majority of recurrent clinical aspects at the outset of ailment were [67-69], along with that Fever at low level not above 100.4°F/38°C in 99 percent cases. In several patients other common features were fatigue, dry cough, anorexia, dyspnea etc [70-72]. However, the fact has to be understood that fever might not be a universal finding.

10. Outline and Guidance

Recently December 2019, a unique/mutant virus, identified as SARS-CoV-2, basically recognized as the significant reason for onset of acute respiratory disease first at Wuhan, a province of China. Nevertheless all began from there; reports have revealed the outbreak of first COVID-19 disease which has expanded widely greater than eighty thousand cases in China itself and thereby increasing cases worldwide. Hence WHO has declared a general well being extremity recently in January 2020. Later it has been characterized as a epidemic in March 2020. The symptoms of COVID-19

displayed fever and persistent dry cough in patients or respiratory tract symptoms. Those infected asymptomatic patients who have had recent close contact also confirmed with COVID-19. Since these asymptomatic patients travelled to areas where community transmission has been reported (eg, China, South Korea, most of Europe [including Italy], Iran, Japan and several other parts of the world including India. Every clinician has to be conscious about the threat of COVID-19 in cases with severe respiratory diseases and take effective precautionary measures while treating those patients. Any suspects of COVID-19 infection have to be notified through helpline and promptly control measures have to be implemented. The best precautionary, measure to decrease the menace for transference of infection to community by cleaning hands daily, follow personal hygiene (eg, cover mouth while cough), and keep away from mass gatherings as well as avoid local association with infective persons. Social distancing from each individual is counselled, especially in areas which already been community transmitted.

Author Contributions

Affiliation

Department of Biotechnology, Sri Padmavati Mahila Visvavidyalayam (Women's) University, Andhra Pradesh, Tirupati-India.

Contributions

KRP and KRD contributed in writing, drawing figures and tables in this review article. KRP solely drafted this review article.

Compliance with Ethics Requirements

NIL

ACKNOWLEDGEMENT

KRP is thankful to the Department of Biotechnology, Sri Padmavati Mahila Visva Vidyalyayam (Women's) University, Tiruapti-India.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

References

1. Centers for Disease Control and Prevention: coronavirus disease 2019 (COVID-19) – situationsummary. (2020). Accessed: March 11, 2020: <https://www.cdc.gov/coronavirus/2019-ncov/summary.html>.
2. Centers for Disease Control and Prevention: human coronavirus types. (2020). Accessed: March 11, 2020: <https://www.cdc.gov/coronavirus/types.html>.
3. WHO: coronavirus disease (COVID-2019) situation reports. (2020). Accessed: March 11, 2020: <https://www.who.int/emergencies/diseases/novel-coronavirus2019/situation-reports>.
4. Perlman S, Netland J. Coronaviruses post-SARS: update on replication and pathogenesis. *Nature Reviews Microbiology*. (2009); 7(6):439–50. <https://doi.org/10.1038/nrmicro2147> WOS:000266451100012. PMID: 19430490.
5. Spaan W, Cavanagh D, Horzinek MC. Coronaviruses: structure and genome expression. *J Gen Virol*. (1988); 69 (Pt 12):2939–52. Epub 1988/12/01. <https://doi.org/10.1099/0022-1317-69-12-2939> PMID: 3058868.
6. Gonzaaez JM, Gomez-Puertas P, Cavanagh D, Gorbalenya AE, Enjuanes L. A comparative

- sequence analysis to revise the current taxonomy of the family Coronaviridae. *Archives of Virology*. (2003); 148 (11):2207–35. <https://doi.org/10.1007/s00705-003-0162-1> WOS:000186399900009. PMID: 14579179.
7. Weiss, S.R.; Navas-Martin, S. Coronavirus pathogenesis and the emerging pathogen severe acute respiratory syndrome coronavirus. *Microbiol. Mol. Biol. Rev.* (2005), 69, 635–664.
 8. Abraham S, Kienzle TE, Lapps W, Brian DA. Deduced sequence of the bovine coronavirus spike protein and identification of the internal proteolytic cleavage site. *Virology*. (1990); 176(1):296–301.
 9. Luytjes W, Sturman LS, Bredenbeek PJ, Charite J, van der Zeijst BA, Horzinek MC, Spaan WJ. Primary structure of the glycoprotein E2 of coronavirus MHV-A59 and identification of the trypsin cleavage site. *Virology*. (1987); 161(2):479–487.
 10. De Groot RJ, Luytjes W, Horzinek MC, van der Zeijst BA, Spaan WJ, Lenstra JA. Evidence for a coiled-coil structure in the spike proteins of coronaviruses. *J Mol Biol.* (1987); 196(4):963–966.
 11. Armstrong J, Niemann H, Smeekens S, Rottier P, Warren G. Sequence and topology of a model intracellular membrane protein, E1 glycoprotein, from a coronavirus. *Nature*. (1984); 308(5961): 751–752.
 12. Nal B, Chan C, Kien F, Siu L, Tse J, Chu K, Kam J, Staropoli I, Crescenzo-Chaigne B, Escriou N, van der Werf S, Yuen KY, Altmeyer R. Differential maturation and subcellular localization of severe acute respiratory syndrome coronavirus surface proteins S, M and E. *The Journal of general virology*. (2005); 86(Pt 5):1423–1434.
 13. Graham, R.L.; Donaldson, E.F.; Baric, R.S. A decade after SARS: Strategies for controlling emerging coronaviruses. *Nat. Rev. Microbiol.* (2013), 11, 836–848.
 14. Frieman, M.; Baric, R. Mechanisms of Severe Acute Respiratory Syndrome Pathogenesis and Innate Immunomodulation. *Microbiol. Mol. Biol. Rev. MMBR* (2008), 72, 672–685.
 15. Peiris, J.S.M.; Guan, Y.; Yuen, K.Y. Severe acute respiratory syndrome. *Nat. Med.* (2004), 10, S88–S97.
 16. Wang, M.; Yan, M.; Xu, H.; Liang, W.; Kan, B.; Zheng, B.; Chen, H.; Zheng, H.; Xu, Y.; Zhang, E.; et al. SARS-CoV Infection in a Restaurant from Palm Civet. *Emerg. Infect. Dis.* (2005), 11, 1860–1865.
 17. Hu, B.; Ge, X.; Wang, L.-F.; Shi, Z. Bat origin of human coronaviruses. *Viol. J.* (2015), 12, 221.
 18. Kim, Y.; Cheon, S.; Min, C.-K.; Sohn, K.M.; Kang, Y.J.; Cha, Y.-J.; Kang, J.I.; Han, S.K.; Ha, N.Y.; Kim, G.; et al. Spread of Mutant Middle East Respiratory Syndrome Coronavirus with Reduced Affinity to Human CD26 during the South Korean Outbreak. *mBio* (2016), 7.
 19. Oboho, I.K.; Tomczyk, S.M.; Al-Asmari, A.M.; Banjar, A.A.; Al-Mugti, H.; Aloraini, M.S.; Alkhalidi, K.Z.; Almohammadi, E.L.; Alraddadi, B.M.; Gerber, S.I.; et al. 2014 MERS-CoV Outbreak in Jeddah—A Link to Health Care Facilities. *N. Engl. J. Med.* (2015), 372, 846–854.
 20. Neuman BW, Kiss G, Kunding AH, Bhella D, Baksh MF, Connelly S, Droese B, Klaus JP, Makino S, Sawicki SG, Siddell SG, Stamou DG, Wilson IA, Kuhn P, Buchmeier MJ. A structural analysis of M protein in coronavirus assembly

- and morphology. *Journal of structural biology*. 2011; 174(1):11–22.
21. Enjuanes, L.; Almazan, F.; Sola, I.; Zuniga, S. Biochemical aspects of coronavirus replication and virus-host interaction. *Annu. Rev. Microbiol.* 2006, 60, 211–230.
 22. Bosch, B.J.; van der Zee, R.; de Haan, C.A.; Rottier, P.J. The coronavirus spike protein is a class I virus fusion protein: Structural and functional characterization of the fusion core complex. *J. Virol.* 2003, 77, 8801–8811.
 23. White, J.M.; Delos, S.E.; Brecher, M.; Schornberg, K. Structures and mechanisms of viral membrane fusion proteins: Multiple variations on a common theme. *Crit. Rev. Biochem. Mol. Biol.* 2008, 43, 189–219.
 24. Xu, Y.; Cole, D.K.; Lou, Z.; Liu, Y.; Qin, L.; Li, X.; Bai, Z.; Yuan, F.; Rao, Z.; Gao, G.F. Construct design, biophysical, and biochemical characterization of the fusion core from mouse hepatitis virus (a coronavirus) spike protein. *Protein Expr. Purif.* 2004, 38, 116–122.
 25. Supekar, V.M.; Bruckmann, C.; Ingallinella, P.; Bianchi, E.; Pessi, A.; Carfi, A. Structure of a proteolytically resistant core from the severe acute respiratory syndrome coronavirus S2 fusion protein. *Proc. Natl. Acad. Sci. U. S. A.* 2004, 101, 17958–17963.
 26. Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, et al. (2003) Coronavirus as a possible cause of severe acute respiratory syndrome. *Lancet* 361: 1319–1325.
 27. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA (2012) Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 367: 1814–1820.
 28. De Haan CA, Rottier PJ (2005) Molecular interactions in the assembly of coronaviruses. *Advances in virus research* 64: 165–230.
 29. Bosch BJ, van der Zee R, de Haan CA, Rottier PJ (2003) The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex. *Journal of virology* 77: 8801–8811.
 30. Inoue Y, Tanaka N, Tanaka Y, Inoue S, Morita K, et al. (2007) Clathrin-dependent entry of severe acute respiratory syndrome coronavirus into target cells expressing ACE2 with the cytoplasmic tail deleted. *Journal of virology* 81: 8722–8729.
 31. Wang H, Yang P, Liu K, Guo F, Zhang Y, et al. (2008) SARS coronavirus entry into host cells through a novel clathrin- and caveolae-independent endocytic pathway. *Cell research* 18: 290–301.
 32. R.J. Hulswit, C.A. de Haan, B.J. Bosch. Coronavirus spike protein and tropism changes. *Adv. Virus Res.*, 96 (2016), pp. 29-57
 33. S.C. Harrison. Viral membrane fusion. *Virology*. 479–480 (2015), pp. 498-507.
 34. Hsue B, Masters PS. A bulged stem-loop structure in the 3' untranslated region of the genome of the coronavirus mouse hepatitis virus is essential for replication. *Journal of virology*. 1997; 71(10): 7567–7578.
 35. Hsue B, Hartshorne T, Masters PS. Characterization of an essential RNA secondary structure in the 3' untranslated region of the murine coronavirus genome. *Journal of virology*. 2000; 74(15):6911– 6921.
 36. Sawicki SG, Sawicki DL, Siddell SG. A contemporary view of coronavirus transcription. *Journal of virology*. 2007; 81(1):20–29.

37. Bentley K, Keep SM, Armesto M, Britton P. Identification of a noncanonically transcribed subgenomic mRNA of infectious bronchitis virus and other gammacoronaviruses. *Journal of virology*. 2013; 87(4):2128–2136.
38. Keck JG, Makino S, Soe LH, Fleming JO, Stohlman SA, Lai MM. RNA recombination of coronavirus. *Advances in experimental medicine and biology*. 1987; 218:99–107.
39. Chang, Y.-J.; Liu, C.Y.-Y.; Chiang, B.-L.; Chao, Y.-C.; Chen, C.-C. Induction of IL-8 Release in Lung Cells via Activator Protein-1 by Recombinant Baculovirus Displaying Severe Acute Respiratory Syndrome-Coronavirus Spike Proteins: Identification of Two Functional Regions. *J. Immunol.* 2004, 173, 7602–7614. [CrossRef] [PubMed]
40. 187. Li, S.-W.; Lai, C.-C.; Ping, J.-F.; Tsai, F.-J.; Wan, L.; Lin, Y.-J.; Kung, S.H.; Lin, C.W. Severe acute respiratory syndrome coronavirus papain-like protease suppressed alpha interferon-induced responses through downregulation of extracellular signal-regulated kinase 1-mediated signalling pathways. *J. Gen. Virol.* 2011, 92, 1127–1140. [CrossRef] [PubMed]
41. 188. Varshney, B.; Lal, S. SARS-CoV Accessory Protein 3b Induces AP-1 Transcriptional Activity through Activation of JNK and ERK Pathways. *Biochemistry* 2011, 50, 5419–5425. [CrossRef] [PubMed]
42. Kindrachuk, J.; Ork, B.; Hart, B.J.; Mazur, S.; Holbrook, M.R.; Frieman, M.B.; Traynor, D.; Johnson, R.F.; Dyal, J.; Kuhn, J.H.; et al. Antiviral Potential of ERK/MAPK and PI3K/AKT/mTOR Signaling Modulation for Middle East Respiratory Syndrome Coronavirus Infection as Identified by Temporal Kinome Analysis. *Antimicrob. Agents Chemother.* 2015, 59, 1088–1099.
43. Kono, M.; Tatsumi, K.; Imai, A.M.; Saito, K.; Kuriyama, T.; Shirasawa, H. Inhibition of human coronavirus 229E infection in human epithelial lung cells (L132) by chloroquine: Involvement of p38 MAPK and ERK. *Antivir. Res.* 2008, 77, 150–152.
44. Mizutani, T.; Fukushi, S.; Saijo, M.; Kurane, I.; Morikawa, S. JNK and PI3k/Akt signaling pathways are required for establishing persistent SARS-CoV infection in Vero E6 cells. *Biochim. Biophys. Acta* 2005, 1741, 4–10.
45. Kanzawa, N.; Nishigaki, K.; Hayashi, T.; Ishii, Y.; Furukawa, S.; Niino, A.; Yasui, F.; Kohara, M.; Morita, K.; Matsushima, K.; et al. Augmentation of chemokine production by severe acute respiratory syndrome coronavirus 3a/X1 and 7a/X4 proteins through NF- κ B activation. *FEBS Lett.* 2006, 580, 6807–6812.
46. Liu, M.; Yang, Y.; Gu, C.; Yue, Y.; Wu, K.K.; Wu, J.; Zhu, Y. Spike protein of SARS-CoV stimulates cyclooxygenase-2 expression via both calcium-dependent and calcium-independent protein kinase C pathways. *FASEB J.* 2007, 21, 1586–1596.
47. Fung, T.S. Molecular Characterization of Cellular Stress Responses during Coronavirus Infection. Ph.D. Thesis, Nanyang Technological University, Singapore, Singapore, 2015.
48. Kopecky-Bromberg, S.A.; Martinez-Sobrido, L.; Palese, P. 7a Protein of Severe Acute Respiratory Syndrome Coronavirus Inhibits Cellular Protein Synthesis and Activates p38 Mitogen-Activated Protein Kinase. *J. Virol.* 2006, 80, 785–793.
49. Tan, Y.-J.; Teng, E.; Shen, S.; Tan, T.H.P.; Goh, P.-Y.; Fielding, B.C.; Ooi, E.E.; Tan, H.C.;

- Lim, S.G.; Hong, W. A Novel Severe Acute Respiratory Syndrome Coronavirus Protein, U274, Is Transported to the Cell Surface and Undergoes Endocytosis. *J. Virol.* 2004, 78, 6723–6734.
50. Jimenez-Guardeño, J.M.; Nieto-Torres, J.L.; DeDiego, M.L.; Regla-Nava, J.A.; Fernandez-Delgado, R.; Castaño-Rodriguez, C.; Enjuanes, L. The PDZ-Binding Motif of Severe Acute Respiratory Syndrome Coronavirus Envelope Protein Is a Determinant of Viral Pathogenesis. *PLoS Pathog.* 2014, 10, e1004320.
 51. Lee, C.-H.; Chen, R.-F.; Liu, J.-W.; Yeh, W.-T.; Chang, J.-C.; Liu, P.-M.; Eng, H.L.; Lin, M.C.; Yang, K.D. Altered p38 Mitogen-Activated Protein Kinase Expression in Different Leukocytes with Increment of Immunosuppressive Mediators in Patients with Severe Acute Respiratory Syndrome. *J. Immunol.* 2004, 172, 7841–7847.
 52. Chen N, Zhou M, Dong X, et al.: Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020, 395:507-13. [10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7)
 53. Letko M, Marzi A, Munster V: Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol.* 2020, Accessed: March 15, 2020: <https://www.nature.com/articles/s41564-020-0688-y>. [10.1038/s41564-020-0688-y](https://doi.org/10.1038/s41564-020-0688-y)
 54. Gralinski LE, Menachery VD: Return of the Coronavirus: 2019-nCoV. *Viruses.* 2020, 12:E135. Accessed: March 11, 2020: <https://www.mdpi.com/1999-4915/12/2/135/htm>. [10.3390/v12020135](https://doi.org/10.3390/v12020135)
 55. Evidence of recombination in coronaviruses implicating pangolin origins of nCoV-2019.(2020).Accessed:March11,2020: <https://www.biorxiv.org/content/10.1101/2020.02.07.939207v1>.
 56. Peng X, Xu X, Li Y, Cheng L, Zhou X, Ren B: Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci.* 2020, 12:9. [10.1038/s41368-020-0075-9](https://doi.org/10.1038/s41368-020-0075-9)
 57. Identification of a pangolin niche for a 2019-nCoV-like coronavirus through an extensive meta-metagenomic search. (2020). Accessed: March 11, 2020: <https://www.biorxiv.org/content/10.1101/2020.02.08.939660v2>.
 58. Centers for Disease Control and Prevention: how COVID-19 spreads. (2020). Accessed:March11,2020: <https://www.cdc.gov/coronavirus/2019ncov/about/transmission.html>.
 59. Bai Y, Yao L, Wei T, et al. Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA* 2020.
 60. Hu Z, Song C, Xu C, et al. Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020.
 61. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020.
 62. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020; 395:514.
 63. Liu W, Zhang Q, Chen J, et al. Detection of Covid-19 in Children in Early January 2020 in Wuhan, China. *N Engl J Med* 2020.

64. Liu YC, Liao CH, Chang CF, et al. A Locally Transmitted Case of SARS-CoV-2 Infection in Taiwan. N Engl J Med 2020; 382:1070.
65. Wei M, Yuan J, Liu Y, et al. Novel Coronavirus Infection in Hospitalized Infants Under 1 Year of Age in China. JAMA 2020.
66. World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report-28.https://www.who.int/docs/default-source/coronaviruse/situationreports/20200217-sitrep-28-covid-19.pdf?sfvrsn=a19cf2ad_2 (Accessed on February 18, 2020).
67. Japanese National Institute of Infectious Diseases. Field Briefing: Diamond Princess COVID-19 Cases, 20 Feb Update. <https://www.niid.go.jp/niid/en/2019-ncov-e/9417-covid-dp-fe-02.html> (Accessed on March 01, 2020).
68. Wang Y, Liu Y, Liu L, et al. Clinical outcome of 55 asymptomatic cases at the time of hospital admission infected with SARS-Coronavirus-2 in Shenzhen, China. J Infect Dis 2020.
69. Pan F, Ye T, Sun P, et al. Time Course of Lung Changes On Chest CT During Recovery From 2019 Novel Coronavirus (COVID-19) Pneumonia. Radiology 2020; :200370.
70. Shi H, Han X, Jiang N, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. Lancet Infect Dis 2020.
71. Chang, Lin M, Wei L, et al. Epidemiologic and Clinical Characteristics of Novel Coronavirus Infections Involving 13 Patients Outside Wuhan, China. JAMA 2020.
72. Xu XW, Wu XX, Jiang XG, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. BMJ 2020; 368:m606.
